

Anaphalioides trinervis

SYNONYMS

Helichrysum micranthum DC., *Anaphalis keriensis* (A.Cunn.) C.J.Webb;
Anaphalis trinervis (A.Cunn.) F.Muell.; *Gnaphalium keriense* A.Cunn.;
Anaphalioides keriensis (A.Cunn.) Kirp.; *Gnaphalium novae-zelandiae*
Sch.Bip; *Gnaphalium lyallii* Hook.f.; *Anaphalis trinervis* var. *lyallii* (Hook.f.)
F.Muell.; *Gnaphalium adhaerens* Col.

FAMILY

Asteraceae

AUTHORITY

Anaphalioides trinervis (G.Forst.) Anderb.

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Herbs - Dicotyledonous composites

NVS CODE

ANATRI

CHROMOSOME NUMBER

2n = 28

CURRENT CONSERVATION STATUS

2017 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

FLOWER COLOURS

White, Yellow

LIFE CYCLE

Pappate cypselae dispersed by wind (Thorsen et al., 2009).

ETYMOLOGY

anaphalioides: Resembling the genus *Anaphalis*, the derivation of which is unclear

REFERENCES AND FURTHER READING

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 2009 Vol. 11 No. 4 pp. 285-309

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/anaphalioides-trinervis/>



Kauaeranga Valley, December. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Remutaka Rail Trail. Photographer: Jeremy R. Rolfe, Date taken: 02/12/2006, Licence: CC BY.