

Anisotome imbricata var. imbricata

FAMILY

Apiaceae

AUTHORITY

Anisotome imbricata (Hook.f.) Cockayne var. imbricata

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

CURRENT CONSERVATION STATUS

2017 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

WETLAND PLANT INDICATOR STATUS RATING

FACU: Facultative Upland

Occasionally is a hydrophyte but usually occurs in uplands (non-wetlands).

FLOWER COLOURS

Yellow

LIFE CYCLE

Winged mericarps are dispersed by wind (Thorsen et al., 2009).

ETYMOLOGY

anisotome: Unequal sided

imbricata: From the Latin imbricatus 'tiled', refers to overlapping features such as leaves

REFERENCES AND FURTHER READING

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309.

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/anisotome-imbricata-var-imbricata/>



Old Woman Range. Photographer: John Barkla, Licence: CC BY.



Harris Mts. Photographer: John Barkla, Licence: CC BY.