

Celmisia incana

COMMON NAMES

white mountain daisy

FAMILY

Asteraceae

AUTHORITY

Celmisia incana Hook.f.

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Herbs - Dicotyledonous composites

NVS CODE

CELINC

CHROMOSOME NUMBER

2n = 108

CURRENT CONSERVATION STATUS

2017 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

DISTRIBUTION


Endemic. North and South Islands from Te Moehau (Coromandel Peninsula) south to Otago

HABITAT

Montane to alpine in grassland, herbfield, boulderfield, on rock outcrops and tors and other similar rocky places.

DETAILED DESCRIPTION

Stems stout, woody, up to ± 100 mm diameter; branches stout, woody, clad in long-persistent reflexed leaves; living leaves in close rosettes, patent. Lamina 20–40 × 10–15 mm, obovate-oblong, coriaceous; upper surface ± densely clad in appressed white tomentum forming a pellicle; lower surface densely clad in similar but more appressed tomentum, midrib evident to obscured; apex subacute to obtuse, often apiculate; margins very slightly recurved, remotely denticulate, narrowed to very short petiole up to 5 mm wide, or sometimes directly into thin almost glabrous striate sheath c.10–15 × 7–10 mm. Scape slender, up to 120 mm long, often short at flowering stage, densely clad in floccose hairs. Capitula 25–35 mm diameter; involucre bracts linear-subulate, many, 10–15 mm long, floccose on outer surface, glandular near apex. Ray-florets narrow, up to 12 mm long, white; limb gradually widening to apex. Disk-florets funnelform, c.7–8 mm. long, teeth narrow-triangular; anthers usually distinctly but shortly tailed. Achenes 3.0–3.5 mm long, compressed-cylindric; ribs rather obscure, clad in rather stiff ascending hairs. Pappus of white or sordid-white slender, minutely barbellate hairs up to 7–8 mm long.



Lyell Range. Photographer: Jane Gosden, Date taken: 15/04/2015, Licence: CC BY-NC-SA.



Mount Ruapehu. Photographer: Jeremy R. Rolfe, Date taken: 08/02/2012, Licence: CC BY.

SIMILAR TAXA

Easily recognised by the silvery white leaves which are hairy on both sides. It is most likely to be confused with *Celmisia hectorii* which has also has silvery white hairy leaves but in that species the leaves are linear-spathulate, linear-oblong to linear-obovate instead of obovate-oblong. *Celmisia allanii* regarded by some New Zealand botanists as distinct is seen here as part of the natural variation of *C. incana*. From *C. incana* it is said to differ by having smaller leaves, and by the tomentum of the underleaves being floccose rather than satiny but the distinctions are not clear cut.

FLOWERING

September - March

FLOWER COLOURS

White, Yellow

FRUITING

November - May

LIFE CYCLE

Pappate cypselae are dispersed by wind (Thorsen et al., 2009).

PROPAGATION TECHNIQUE

Best grown in non-humid climates. *Celmisia incana* is one of the few *Celmisia* that generally grows well in most garden conditions. However, it can be fickle. Best grown from fresh seed and planted in a fertile, free draining semi-shaded situation. Dislikes excessive moisture, and humidity.

ETYMOLOGY

celmisia: Apparently named after Kelmis, one of Idaean Dactyls, a group of skilled mythical beings associated with the Mother Goddess Rhea in Greek mythology. Kelmis, whose name means 'casting', was a blacksmith and childhood friend of Zeus, son of Rhea and later king of the gods. In Ovid's 'Metamorphoses', Kelmis is described as offending Zeus who turned him into adamant so he was as hard as a tempered blade

incana: Hoary (greyish white haired)

TAXONOMIC NOTES

Celmisia allanii W.Martin is sometimes included within *C. incana* by New Zealand botanists. But this informal view has yet to be properly tested taxonomically. For this reason *C. allanii* is regarded as distinct from *C. incana*.

CULTIVATION

Occasionally available from specialist native plant nurseries.

ATTRIBUTION

P.J. de Lange (7 April 2009). Description adapted from Allan (1961)

Some of this factsheet information is derived from [Flora of New Zealand Online](#) and is used under a [Creative Commons Attribution 3.0 New Zealand licence](#).

REFERENCES AND FURTHER READING

Allan, H.H. 1961: Flora of New Zealand. Vol. I, Government Printer, Wellington.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309

NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Celmisia incana* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network. <https://www.nzpcn.org.nz/flora/species/celmisia-incana/> (Date website was queried)

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/celmisia-incana/>

PDF DATE

25 September 2024