

# Coprosma propinqua var. propinqua

## COMMON NAMES

mingimingi

## FAMILY

Rubiaceae

## AUTHORITY

*Coprosma propinqua* A.Cunn. var. *propinqua*

## FLORA CATEGORY

Vascular – Native

## ENDEMIC TAXON

Yes

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## STRUCTURAL CLASS

Trees & Shrubs - Dicotyledons

## NVS CODE

COPPVP

## CHROMOSOME NUMBER

2n = 44

## CURRENT CONSERVATION STATUS

2017 | Not Threatened

## PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

## SIMPLIFIED DESCRIPTION

Very common bushy shrub (or low-growing mound in some coastal areas) with wide-angled branches bearing clusters of pairs of variably shaped dark green glossy narrow leaves. Young leaves with dark stalk. Adult leaves often curved sideways, 10-4mm long by 2-3mm wide, paler underneath and with 1-3 pits. Fruit pale blue.

## WETLAND PLANT INDICATOR STATUS RATING

FAC: Facultative

Commonly occurs as either a hydrophyte or non-hydrophyte (non-wetlands).

## MANAAKI WHENUA ONLINE INTERACTIVE KEY

[Key to \*Coprosma\* species of New Zealand](#)

## FLOWER COLOURS

Green

## LIFE CYCLE

Fleshy drupes are dispersed by frugivory (Thorsen et al., 2009).



Eastbourne. Photographer: Jeremy R. Rolfe, Date taken: 08/06/2001, Licence: CC BY.



Pauatahanui Inlet. Photographer: Jeremy R. Rolfe, Date taken: 08/04/2006, Licence: CC BY.

## ETYMOLOGY

**coprosma:** From the Greek kopros 'dung' and osme 'smell', referring to the foul smell of the species, literally 'dung smell'

**propinqua:** From the Latin propinquus 'near, neighbouring', meaning closely related to another species

## REFERENCES AND FURTHER READING

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

## MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/coprosma-propinqua-var-propinqua/>

## PDF DATE

17 September 2024