

# Hiya distans

## SYNONYMS

*Hypolepis distans* Hook.

## FAMILY

Dennstaedtiaceae

## AUTHORITY

*Hiya distans* (Hook.) Brownsey & Perrie

## FLORA CATEGORY

Vascular – Native

## ENDEMIC TAXON

No

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## STRUCTURAL CLASS

Ferns

## NVS CODE

HYPDIS

## CHROMOSOME NUMBER

$2n = 56$

## CURRENT CONSERVATION STATUS

2017 | Not Threatened | Qualifiers: TO

## PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

## DISTRIBUTION

Indigenous. New Zealand: North, South (Marlborough, North-West Nelson and North Westland with one record from Banks Peninsula) and Chatham Islands. Also Australia (Tasmania and some Bass Strait Islands). Sparingly naturalised on Norfolk Island (de Lange et al. 2005)

## HABITAT

*Hiya distans* is usually found in swampy areas, especially on peat, or in soils with a deep humus layer. In forest situations (or disturbed forest margins) it often colonises decomposing logs. Very occasionally it is found as a weed of garden centres and nurseries, plants often arising from the peat that is bought into these areas to use as potting mix.

## WETLAND PLANT INDICATOR STATUS RATING

FAC: Facultative

Commonly occurs as either a hydrophyte or non-hydrophyte (non-wetlands).



Kennedy Bay, August. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Kennedy Bay, August. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.

## DETAILED DESCRIPTION

**Rhizome** long-creeping, 1.0–2.25 mm diam., abundantly covered in dark red-brown hairs 2.0–3.5 mm long, giving rise to stipes at intervals of 30–70 mm. **Stipes** 50–500–(600) mm long, 0.75–1.25 mm diam., dark red-brown, bearing a few scattered red-brown hairs at base. **Laminae** narrowly elliptic to elliptic or ovate, (150)–250–600–(950) × 90–300–(400) mm, rather stiff and harsh to the touch, bipinnate at apex to almost tripinnate at base (truly tripinnate in largest specimens). **Rachis** dark red-brown becoming paler only at the very apex, bearing scattered pale brown hairs. **Primary pinnae** in 20–40 pairs (more in very large specimens), opposite or subopposite arising at c. 90° to rachis, the longest at or below middle 50–200 × 17–60 mm; lowest pinnae (20)–40–120–(200) mm apart, middle pinnae (10)–20–50–(60) mm apart, narrowly ovate or triangular, midrib narrowly winged throughout; lowest pair orientated at 90° to plane of frond in largest specimens. **Secondary pinnae** sessile or shortly stalked, ovate to narrowly ovate, longest 8–30–(40) × 4–7–(10) mm, doubly serrate with 3–9 primary serrations on each side of longest secondary pinna (divided to the midrib in tripinnate forms); those on lower primary pinnae ± equal in length, only decreasing in size towards the very apex of each pinna. **Veins** reaching margin at a sinus (emarginate). **Hairs** virtually absent on upper surface of frond, a few scattered pale-brown non-glandular hairs on underside of costae and midribs. **Sori** in 2–8 pairs on longest secondary pinnae, marginal, terminating veins on sides of ultimate segments, protected by reflexed, incised membranous flaps, dark brown, ± smooth.

## SIMILAR TAXA

None.

## FLOWERING

Not applicable—spore producing

## FLOWER COLOURS

No flowers

## FRUITING

Not applicable—spore producing

## PROPAGATION TECHNIQUE

Easily grown from fresh spores and by division of established plants. Should be planted in a humus rich (peat is ideal), damp soil. Does best in a warm sunny place but once established tolerant of a range of situations of moisture levels.

## ETYMOLOGY

**distans**: Distant (widely spaced female flowers)

## ATTRIBUTION

Fact sheet prepared for NZPCN by P.J. de Lange (8 November 2012). Description from Brownsey & Chinnock (1984).

## REFERENCES AND FURTHER READING

Brownsey PJ, Chinnock RJ. 1984. A Taxonomic revision of the New Zealand species of *Hypolepis*. *New Zealand Journal of Botany* 22(1): 43–80. <https://doi.org/10.1080/0028825X.1984.10425234>.

de Lange PJ, Gardner RO, Sykes WR, Crowcroft GM, Cameron EK, Stalker F, Christian ML, Braggins JE. 2005. Vascular flora of Norfolk Island: some additions and taxonomic notes. *New Zealand Journal of Botany* 43(4): 563–596. <https://doi.org/10.1080/0028825X.2005.9512975>.

Perrie LR, Shepherd LD, Brownsey PJ. 2018. *Hiya distans* (Dennstaedtiaceae), a new combination for an Australasian fern previously classified in *Hypolepis*. *New Zealand Journal of Botany* 56(4): 406–414. <https://doi.org/10.1080/0028825X.2018.1526807>.

## NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Hiya distans* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network. <https://www.nzpcn.org.nz/flora/species/hiya-distans/> (Date website was queried)

## MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/hiya-distans/>

**PDF DATE**

17 September 2024