

Pomaderris edgerleyi

BIOSTATUS

Native

CATEGORY

Vascular

STRUCTURAL CLASS

Trees & Shrubs - Dicotyledons

SIMPLIFIED DESCRIPTION

Low-growing shrub to 1m tall with rusty fuzzy twigs bearing small wrinkled leathery oval leaves with small teeth along the edge and clusters of yellowish flowers inhabiting Northland. Leaves 15–25 mm long by 7–10 mm wide, fuzzy underneath. Fruit dry, small.

FLOWER COLOURS

Cream, Yellow

DISTRIBUTION

Endemic. New Zealand: North Island (Te Paki south to about Dargaville).

CURRENT CONSERVATION STATUS

2023 | At Risk – Declining | Qualifiers: DPS, DPT, PF

DETAILED TAXONOMY

FAMILY

Rhamnaceae

AUTHORITY

Pomaderris edgerleyi Hook.f.

SYNONYMS

Pomaderris prunifolia var. *edgerleyi* (Hook.f.) L.B.Moore

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

OTHER INFORMATION

CULTIVATION

Occasionally sold by specialist native plant nurseries

ETYMOLOGY

pomaderris: Lid skin

edgerleyi: Named after John Edgerley (1814 - 1849) who spent time in New Zealand and Australia

NVS CODE

POMEDG

CHROMOSOME NUMBER

2n = 36, 37



Hikurua / de Surville Cliffs, Te Paki.
Photographer: Jeremy R. Rolfe, Date taken:
19/10/2009, Licence: CC BY.



Hikurua / de Surville Cliffs, Te Paki.
Photographer: Jeremy R. Rolfe, Date taken:
26/02/2011, Licence: CC BY.

PREVIOUS CONSERVATION STATUSES

2017 | At Risk – Declining | Qualifiers: DP

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/pomaderris-edgerleyi/>

PDF DATE

15 October 2024