

Tanacetum parthenium

COMMON NAMES

feverfew

FAMILY

Asteraceae

AUTHORITY

Tanacetum parthenium (L.) Sch.Bip.

FLORA CATEGORY

Vascular – Exotic

STRUCTURAL CLASS

Herbs - Dicotyledonous composites

NVS CODE

TANPAR

CONSERVATION STATUS

Not applicable

HABITAT

Dry soils, waste places, road sides. Introduced from Europe, feverfew is now a common garden plant. It can be cultivated in all parts of New Zealand.

FLOWER COLOURS

White, Yellow

LIFE CYCLE

Perennial

YEAR NATURALISED

1875

ORIGIN

Balkan Peninsula

REASON FOR INTRODUCTION

Herbal remedy, ornamental

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/tanacetum-parthenium/>

PDF DATE

17 September 2024



Double-flowered form. Hutt River north of Stokes Valley, Lower Hutt. Photographer: Jeremy R. Rolfe, Date taken: 27/01/2007, Licence: CC BY.



Double-flowered form. Hutt River north of Stokes Valley, Lower Hutt. Photographer: Jeremy R. Rolfe, Date taken: 27/01/2007, Licence: CC BY.

